

## POLICY SECTIONS

[POLICY DESCRIPTION](#) | [DISCLAIMER](#) | [RELATED POLICIES](#) | [INDICATIONS AND/OR LIMITATIONS OF COVERAGE](#) | [EXCLUSION CRITERIA](#) | [MEDICATION MANAGEMENT](#) | [ATTACHMENTS](#) | [APPLICABLE CPT / HCPCS PROCEDURE CODES](#) | [APPROVAL HISTORY](#) | [REFERENCES](#) | [APPENDIX](#)

## DISCLAIMER

This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. Policies are not a supplementation or recommendation for treatment; Providers are solely responsible for the diagnosis, treatment and clinical recommendations for the Member. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (e.g., will be paid for by Molina) for a particular Member. The Member's benefit plan determines coverage – each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their Providers will need to consult the Member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid Members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this MCP and provide the directive for all Medicare members. References included were accurate at the time of policy approval and publication.

## POLICY DESCRIPTION

To define and describe the accepted indications for Abecma (idecabtagene vicleucel) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

## INDICATIONS and/or LIMITATIONS OF COVERAGE

### A. PREFERRED MEDICATION GUIDANCE FOR INITIAL REQUEST:

1. When health plan Medicaid coverage provisions—including any applicable PDLs (Preferred Drug Lists)—conflict with the coverage provisions in this drug policy, health plan Medicaid coverage provisions take precedence per the **Preferred Drug Guidelines**; **OR**
2. When health plan Exchange coverage provisions—including any applicable PDLs (Preferred Drug Lists)—conflict with the coverage provisions in this drug policy, health plan Exchange coverage provisions take precedence per the **Preferred Drug Guidelines**; **OR**
3. When Health Plans utilize NCH UM Oncology Clinical Policies as the initial clinical criteria, and there is no Health Plan PDL applicable, the Preferred Drug Guidelines shall follow NCH recommended agents/regimens/preferred drugs; **AND**
4. Continuation requests of previously approved, non-preferred medication are not subject to this provision; **AND**
5. When applicable, generic alternatives are preferred over brand-name drugs; **AND**
6. For Health Plans utilizing NCH UM Oncology Clinical Policies, when there is a documented drug shortage, disease progression, contraindication, or confirmed intolerance to a Preferred drug/regimen, per NCH Policy and Pathway, the available alternative product may be used if deemed medically appropriate and the indication is listed in a standard reference compendium or accepted peer review literature. For a list of current drug shortages, please refer to FDA drug shortage website in the reference section.

## B. Multiple Myeloma

1. Abecma (idecabtagene vicleucel) may be used for adult members with relapsed/refractory multiple myeloma that have progressed on 4 or more lines of therapy; **AND**
2. Members must have triple refractory disease as defined in the KaMMA trial: refractory to an immunomodulatory agent (e.g., lenalidomide, thalidomide, pomalidomide), a proteasome inhibitor (e.g., bortezomib, carfilzomib, ixazomib), and an anti-CD38 antibody (e.g., daratumumab, isatuximab); **AND**
3. Members must have measurable disease or evidence of disease progression from the last line of therapy for multiple myeloma.

## EXCLUSION CRITERIA

- A. Disease progression while receiving Abecma (idecabtagene vicleucel).
- B. Concurrent use with other anti-myeloma therapy.
- C. The member does NOT have measurable disease defined as any of the following:
  1. Serum M-protein greater or equal to 1.0 g/dL
  2. Urine M-protein greater or equal to 200 mg/24 h
  3. Serum free light chain (FLC) assay: involved FLC level greater or equal to 10 mg/dL (100 mg/L) provided serum FLC ratio is abnormal.
- D. The member has solitary plasmacytomas or non-secretory myeloma without other evidence of measurable disease.
- E. The member does NOT have adequate bone marrow reserve defined by **ALL** of the following:
  1. Absolute neutrophil count (ANC)  $\geq$  1000/uL
  2. Platelet Count  $\geq$  50,000/uL.
- F. The member does NOT have adequate renal, hepatic, cardiac and pulmonary function defined as:
  1. Creatinine clearance  $>$  45 mL/min
  2. AST and/or ALT  $\leq$  2.5 x ULN and total bilirubin  $\leq$  1.5 x ULN
  3. Cardiac ejection fraction  $\geq$  45%, no evidence of pericardial effusion as determined by an echocardiogram (ECHO), and no clinically significant pleural effusion.
- G. History or presence of CNS disorder.
- H. Prior allogeneic hematopoietic stem cell transplant (HSCT).
- I. Does not exceed duration limit as one time administration.
- J. Dosing exceeds single dose limit of Abecma (idecabtagene vicleucel)  $460 \times 10^6$  CAR-positive T cells.
- K. Investigational use of Abecma (idecabtagene vicleucel) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
  1. Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
  2. Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
  3. Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definition of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of  $<$  0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.

## Molina Clinical Policy

### Abecma™ (idecabtagene vicleucel): Policy Number 403

Last Approval: 8/9/2023

Next Review Due By: August 2024



4. Whether the experimental design, in light of the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
5. That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
6. That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
7. That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

#### MEDICATION MANAGEMENT

- A. Please refer to the FDA label/package insert for details regarding these topics.

#### APPLICABLE CPT / HCPCS PROCEDURE CODES

##### CPT (Current Procedural Terminology) Codes

CPT	Description
0537T	Chimeric antigen receptor T-cell (CAR-T) therapy; harvesting of blood-derived T lymphocytes for development of genetically modified autologous CAR-T cells, per day
0538T	Chimeric antigen receptor T-cell (CAR-T) therapy; preparation of blood-derived T lymphocytes for transportation (e.g., cryopreservation, storage)
0539T	Chimeric antigen receptor T-cell (CAR-T) therapy; receipt and preparation of CAR-T cells for administration
0540T	Chimeric antigen receptor T-cell (CAR-T) therapy; CAR-T cell administration, autologous

##### HCPCS (Healthcare Common Procedure Coding System) Code

HCPCS	Description
Q2055	Idecabtagene vicleucel, up to 460 million autologous B-cell maturation antigen (BCMA) directed CAR-positive T cells, including leukapheresis and dose preparation procedures, per therapeutic dose

**AVAILABLE DOSAGE FORMS:** Single-dose unit infusion bag: frozen suspension of genetically modified autologous T-cells labeled for the specific recipient.

**CODING DISCLAIMER.** Codes listed in this policy are for reference purposes only and may not be all-inclusive. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement. Listing of a service or device code in this policy does not guarantee coverage. Coverage is determined by the benefit document. Molina adheres to Current Procedural Terminology (CPT®), a registered trademark of the American Medical Association (AMA). All CPT codes and descriptions are copyrighted by the AMA; this information is included for informational purposes only. Providers and facilities are expected to utilize industry standard coding practices for all submissions. When improper billing and coding is not followed, Molina has the right to reject/deny the claim and recover claim payment(s). Due to changing industry practices, Molina reserves the right to revise this policy as needed.

#### APPROVAL HISTORY

- 08/09/2023** For multiple myeloma, added indication to clarify that members must have measurable disease or evidence of disease progression from the last line of therapy. Added code Q2055 and removed codes C9399, J3490, J3590, and J9999.
- 08/10/2022** Adopted NCH policy and retired MCP.

#### REFERENCES

# Molina Clinical Policy

## Abecma™ (idecabtagene vicleucel): Policy Number 403

Last Approval: 8/9/2023

Next Review Due By: August 2024



1. Munshi NC, et al. KarMMa Trial. Idecabtagene Vicleucel in Relapsed and Refractory Multiple Myeloma. N Engl J Med. 2021 Feb 25;384(8):705-716.
2. Raje N, et al. CRB-401 Trial. Anti-BCMA CAR T-Cell Therapy bb2121 in Relapsed or Refractory Multiple Myeloma. N Engl J Med. 2019 May 2;380(18):1726-1737.
3. Abecma prescribing information. Celgene Corporation, Summit, NJ 2022.
4. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. J Clin Oncol. 2014 Apr 20;32(12):1277-80.
5. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.
6. Current and Resolved Drug Shortages and Discontinuations Reported to the FDA: <http://www.accessdata.fda.gov/scripts/drugshortages/default.cfm>
7. NCQA UM 2023 Standards and Elements.

### APPENDIX

**Reserved for State specific information.** Information includes, but is not limited to, State contract language, Medicaid criteria and other mandated criteria.