



Request for Prior Authorization

PCSK9 INHIBITORS

(PLEASE PRINT - ACCURACY IS IMPORTANT)

FAX Completed Form To
1 (877) 733-3195
IOWA Provider Services
1 (844) 236-1464

Form with fields for: IA Medicaid Member ID #, Patient name, DOB, Patient address, Provider NPI, Prescriber name, Phone, Prescriber address, Fax, Pharmacy name, Address, Phone, Pharmacy NPI, Pharmacy fax, NDC.

Prior authorization is required for PCSK9 Inhibitors. Payment for a non-preferred PCSK9 Inhibitor will be authorized only for cases in which there is documentation of a previous trial and therapy failure with a preferred agent. Payment will be considered under the following conditions: 1) Patient meets the FDA approved age for indication; and 2) Dosing follows the FDA approved dose for the submitted diagnosis; and 3) Current use of a statin and documentation of adherence to prescribed lipid lowering medications for the previous 90 days is provided (further defined below, by diagnosis); and 4) Is to be prescribed as an adjunct to a low fat diet; and 5) A baseline and current lipid profile is provided. Baseline lipid profile is defined as a lipid profile obtained prior to pharmacologic therapy; and 6) Documentation patient has been counseled on importance of abstinence from tobacco and, if a current smoker, be encouraged to enroll in a smoking cessation program; and 7) The 72-hour emergency supply rule does not apply to PCSK9 Inhibitors. 8) Prescriber and dispensing pharmacy will educate the patient on proper storage and administration. Improperly stored medications will not be replaced. 9) Lost or stolen medication replacement requests will not be authorized. 10) Goal is defined as a 50% reduction in untreated baseline LDL-C. 11) Is prescribed for one of the following diagnoses: Heterozygous Familial Hypercholesterolemia (HeFH), Clinical Atherosclerotic Cardiovascular Disease (ASCVD), Primary Hyperlipidemia (not associated with ASCVD or HeFH), or HoFH. The required trials may be overridden when documented evidence is provided that use of these agents would be medically contraindicated.

Initial requests will be approved for 6 months. Additional requests will be considered under the following conditions: 1) Documentation of positive clinical response to PCSK9 inhibitor therap (current LDL-C lab provided); and 2) Patient continues therapy with a maximally tolerated statin; and 3) Patient has continued compliance with a low-fat diet.

Preferred

[] Praluent [] Repatha

Strength Dosage Instructions Quantity Days Supply

Initial Requests (please see below for renewal requests):

Is patient on a low fat diet: [] Yes [] No

Has patient experienced ≥ 50% reduction in untreated baseline LDL-C with current therapies? [] Yes [] No

Attach baseline (prior to pharmacologic therapy) and current lipid profiles.

Statin to be used as adjunct to PCSK9 inhibitor: _____ Dose: _____

Has patient been counseled on importance of abstinence from tobacco? [] Yes [] No

Request for Prior Authorization

PCSK9 INHIBITORS

(PLEASE PRINT – ACCURACY IS IMPORTANT)

Is patient a current smoker or tobacco user:

Yes No

If yes, has patient been encouraged to enroll in smoking cessation program?

Yes No

Prescriber and dispensing pharmacy will educate patient on proper storage and administration?

Yes No

Heterozygous Familial Hypercholesterolemia (HeFH)

- 1) Total cholesterol > 290mg/dL or LDL-C > 190mg.dL; and
 - a) Presence of tendon xanthomas; or
 - b) In first or second degree relative, one of the following: documented tendon xanthomas, MI at age ≤ 60 years, or total cholesterol > 290mg/dL; or
 - c) Confirmation of diagnosis by gene or receptor testing (attach results); and
- 2) Unable to reach goal LDL-C with a minimum of one high-intensity statin (atorvastatin 40-80 mg or rosuvastatin 20-40 mg) used in combination with ezetimibe 10 mg daily. If patient is unable to tolerate high-intensity statin therapy, a trial with a moderate-intensity statin (e.g., atorvastatin 10-20 mg, rosuvastatin 5-10 mg, pravastatin 40-80 mg, lovastatin 40-80 mg, fluvastatin 80 mg, pitavastatin 1-4 mg, simvastatin 20-40 mg) used in combination with ezetimibe.

Total cholesterol: _____

Date obtained: _____

LDL-C: _____

Date obtained: _____

Presence of tendon xanthomas: Yes No

Any of the following present in first degree relative:

Documented tendon xanthomas MI at age ≤ 60 years Total cholesterol > 290mg/dL

Diagnosis confirmed by gene or receptor testing? Yes (attach results) No

High or Medium- Intensity Statin trial:

Dose: _____ Trial dates: _____

Failure reason: _____

Rationale for medium-intensity statin trial: _____

Plus concurrent ezetimibe trial:

Dose: _____ Trial dates: _____

Failure reason: _____

Medical or contraindication reason to override trial requirements: _____

Clinical Atherosclerotic Cardiovascular Disease (ASCVD)

- 1) History of MI, angina, coronary or other arterial revascularization, stroke, TIA, or PVD of atherosclerotic origin; and
- 2) Unable to reach goal LDL-C with a minimum of one high-intensity statin (atorvastatin 40-80 mg or rosuvastatin 20-40 mg) used in combination with ezetimibe 10 mg daily. If patient is unable to tolerate high-intensity statin therapy, a trial with a moderate-intensity statin (e.g., atorvastatin 10-20 mg, rosuvastatin 5-10 mg, pravastatin 40-80 mg, lovastatin 40-80 mg, fluvastatin 80 mg, pitavastatin 1-4 mg, simvastatin 20-40 mg) used in combination with ezetimibe.

History of any of the following: MI Angina Coronary or other arterial revascularization Stroke TIA PVD of atherosclerotic origin

High or Medium-Intensity Statin trial:

Dose: _____ Trial dates: _____

**Request for Prior Authorization
PCSK9 INHIBITORS**
(PLEASE PRINT – ACCURACY IS IMPORTANT)

Failure reason: _____

Rationale for medium-intensity statin trial: _____

Plus concurrent ezetimibe trial:

Dose: _____ Trial dates: _____

Failure reason: _____

Medical or contraindication reason to override trial requirements: _____

Primary Hyperlipidemia (not associated with ASCVD or HeFH)

- 1) Baseline LDL-C \geq 190 mg/dL; and
- 2) Unable to reach goal LDL-C < 100 mg/dL while on high-intensity statin therapy (atorvastatin 40-80 mg or rosuvastatin 20-40 mg) used in combination with ezetimibe 10 mg daily. If patient is unable to tolerate high-intensity statin therapy, a trial with a moderate-intensity statin (e.g., atorvastatin 10-20 mg, rosuvastatin 5-10 mg, pravastatin 40-80 mg, lovastatin 40-80 mg, fluvastatin 80 mg, pitavastatin 1-4 mg, simvastatin 20-40 mg) used in combination with ezetimibe.

LDL-C: _____ **Date obtained:** _____

High or Medium- Intensity Statin trial:

Dose: _____ Trial dates: _____

Failure reason: _____

Rationale for medium-intensity statin trial: _____

Plus concurrent ezetimibe trial:

Dose: _____ Trial dates: _____

Failure reason: _____

Medical or contraindication reason to override trial requirements: _____

Homozygous Familial Hypercholesterolemia (HoFH)

- 1) Total cholesterol and LDL-C > 600mg/dL and triglycerides within reference range; or
- 2) Confirmation of diagnosis by gene or receptor testing (attach results); and
- 3) Unable to reach goal LDL-C with a minimum of one high-intensity statin (atorvastatin 40-80 mg or rosuvastatin 20-40 mg) used in combination with ezetimibe 10 mg daily. If patient is unable to tolerate high-intensity statin therapy, a trial with a moderate-intensity statin (e.g., atorvastatin 10-20 mg, rosuvastatin 5-10 mg, pravastatin 40-80 mg, lovastatin 40-80 mg, fluvastatin 80 mg, pitavastatin 1-4 mg, simvastatin 20-40 mg) used in combination with ezetimibe.

Total cholesterol: _____ **Date obtained:** _____

LDL-C: _____ **Date obtained:** _____

Triglycerides within reference range? Yes No (attach results)

Diagnosis confirmed by gene or receptor testing? Yes (attach results) No

High or Medium-Intensity Statin trial:

Dose: _____ Trial dates: _____

**Request for Prior Authorization
PCSK9 INHIBITORS**
(PLEASE PRINT – ACCURACY IS IMPORTANT)

Failure reason: _____

Rationale for medium-intensity statin trial: _____

Plus concurrent ezetimibe (Zetia) trial:

Dose: _____ Trial dates: _____

Failure reason: _____

Medical or contraindication reason to override trial requirements: _____

Renewal Requests:

Patient continues therapy with a maximally tolerated statin? Yes No

Current Statin: Drug name: _____ Dose: _____

Patient has continued compliance with a low fat diet? Yes No

Documentation of positive clinical response to PCSK9 Inhibitor therapy (provide current LDL-C lab):

Attach lab results and other documentation as necessary.

Prescriber signature (Must match prescriber listed above.)	Date of submission
--	--------------------

IMPORTANT NOTE: In evaluating requests for prior authorization the consultant will consider the treatment from the standpoint of medical necessity only. If approval of this request is granted, this does not indicate that the member continues to be eligible for Medicaid. It is the responsibility of the provider who initiates the request for prior authorization to establish by inspection of the member's Medicaid eligibility card and, if necessary by contact with the county Department of Human Services, that the member continues to be eligible for Medicaid.